NOAA Climate Science & Services Monthly Climate Update

A look back at July and a preview for Fall (Sept. to Nov.)

Jake Crouch

Climate Scientist, Climate Monitoring Branch, NOAA's National Climatic Data Center

Ed O'Lenic

Seasonal Forecaster, Operations Branch NOAA's Climate Prediction Center


Dr. Jim Angel

State Climatologist
Illinois State Water Survey Prairie Research
Institute & Univ. of Illinois Urbana-Champaign





Global Climate Highlights



July 2012 Precipitation:

- No tropical storms in N. Atlantic Basin.
- Typhoons Vincente & Saola hit SE Asia
- Parts of Argentina have driest July in 50+ years
- Below-average monsoonal rains in India

Dry: Brazil, southern Europe, India, western Aus.

Wet: U.K., China, S. Africa, Queensland Aus.

July 2012 Temperatures:


- 4th warmest July globally (+0.62°C/+1.12°F)
- Land: 3rd warmest; Ocean: 7th warmest
- N. Hemisphere: 2nd warmest July
- January-July: 10th warmest (tied 2001)

Regional Analysis:

- N. America warmer than average
- U.K has coolest July since 2000
- Australia minimum temps 8th coolest on record

Land-Only Precipitation Percent of Normal Jul 2012

(with respect to a 1961-1990 base period)









Global Climate Highlights

- 2nd smallest July Arctic sea ice extent
- Massive ice lost during July: 2.97 million square km
- August 14th: Arctic sea ice among the four lowest summer minimums









U.S. Climate Highlights


Nationally: Warmest Month on Record – 77.6°F


July: National temperatures 0.2°F warmer than July 1936

- Monthly temperatures 3.3°F above average
- Footprint of warm temperatures led to record month

Year-to-Date: First seven months of 2012 record warm

- 33 states record warm for January-July
- Only the Pacific Northwest near-average for the period
- If Aug-Dec near-average, 2012 will be warmest year on record









U.S. Climate Highlights

Nationally: 28th Driest July / 15th Driest Year-to-Date Drought conditions for nearly two-thirds of contiguous United States

July

- Wet in Southwest, Gulf Coast, and Appalachians
- Dry in Central Plains, Midwest, and New England

Year-to-Date

- Wet in Pacific Northwest, Upper Midwest
- Dry in Rockies, Midwest, Mid-South, Northeast


July 2012 Statewide Ranks National Climatic Data Center/NESDIS/NOAA 97 71 73 66 14 101 83 12 97 75 100 100 1 = Driest 118 = Wettest

Near

Normal

January-July 2012 Statewide Ranks

National Climatic Data Center/NESDIS/NOAA







Record

Much

Below

Below


Above

Wettest

Above

Normal

Drought Impacts



http://droughtmonitor.unl.edu/

Released Thursday, August 16, 2012

Author: Michael Brewer/Liz Love-Brotak, NOAA/NESDIS/NCDC






El Niño/La Niña Southern Oscillation (ENSO) Sea Surface Temperature Update

ENSO Neutral Conditions;

Warmer than normal


East Pacific temperatures








1-Month Temperature and Precipitation Forecast








3-Month Temperature and Precipitation Forecast








Latest U.S. Hazards Outlook for Days 3-14









impacts ease

likely

Drought development

by short- and long-range statistical and dynamical forecasts. Short-term events
-- such as individual storms -- cannot be accurately forecast more than a few days in advance.
Use caution for applications -- such as crops -- that can be affected by such events.
"Ongoing" drought areas are approximated from the Drought Monitor (D1 to D4 intensity).
For weekly drought updates, see the latest U.S. Drought Monitor. NOTE: the green improvement areas imply at least a 1-category improvement in the Drought Monitor intensity levels, but do not necessarily imply drought elimination.

Agricultural Impacts

- Estimated reduced yields in corn (13%) and soybeans (12%) -USDA.
- Some corn fields are a total loss, while other fields are cut for silage along with some early harvesting.
- Cooler temperatures across the Midwest and rain in the eastern Midwest in August have helped with soybeans more so than corn.
- Poor pasture, limited hay, and high feed prices, are causing a sell-off of herds.





Secondary Impacts

- Vulnerable water supplies at the farm level – ponds, creeks, shallow wells.
- Problems with low water levels on the lower Mississippi.
- Fire hazard from machinery during harvest.
- Depleted soil moisture this fall could be a problem for establishing winter wheat crop.





For More Information

Today's Presentation:

http://www.ncdc.noaa.gov/sotc/briefings

NOAA's National Climatic Data Center: www.ncdc.noaa.gov

Monthly climate reports (U.S. & Global): www.ncdc.noaa.gov/sotc/

NOAA's Climate Prediction Center: www.cpc.ncep.noaa.gov

U.S. Drought Monitor: http://droughtmonitor.unl.edu/

Climate Portal: www.climate.gov

Illinois State Climatologist: http://www.isws.illinois.edu/atmos/statecli/

Media Contacts

- Brady.Phillips@noaa.gov, 202-407-1298 (NOAA/Comms)
- Katy.Vincent@noaa.gov, 828-257-3136 (NOAA/NCDC)



